
Space Weather Technical Forecast

Issued on Thursday, 04 August 2022 at 00:19 Local

This technical forecast provides a four day assessment of space weather events. The probabilities stated below are for reaching or exceeding the given levels. For more information about space weather impacts please see the Met Office Space Weather Scales <https://www.metoffice.gov.uk/weather/learn-about/space-weather/uk-scales>

Space Weather Forecast Headline: No significant activity.

Analysis of Space Weather Activity over past 24 hours

Solar Activity: Solar activity has been low over the past 24 hours with two C-class flares, the strongest being a C9.2 flare at 03/1708 UTC. Both flares originating from AR3068 Esi/Beta-gamma in the southwest. There is one other region, AR3071 Hsx/alpha, in the southeast. The C9.2 flare was accompanied by a type II radio-burst, indicating a possibility of a CME, however currently there is no evidence on available imagery, further analysis will take place as more imagery becomes available. No other Earth-directed CMEs were observed.

Solar Wind / Geomagnetic Activity: Earth has likely been grazing the High Speed Stream from coronal hole 07, with solar wind speeds, as measured at L1, generally fluctuating between 450 and 550 km/s. Density was around average to just above. The total interplanetary magnetic field (IMF), Bt was generally weak, at below 8 nT. The north-south component, Bz, fluctuated weakly between +6/-5 nT. The Phi angle was negative (towards the Sun). Geomagnetic activity was Quiet (Kp 1-2).

Energetic Particles / Solar Radiation: The high energy proton flux (10 MeV), as measured by GOES-16 was at background levels. The high energy electron flux (greater than 2 MeV) was at background. Electron fluence showed an erratic declining trend, remaining well below the Active threshold (1e8 integrated flux), with an observed fluence value of 4.33e6 integrated pfu recorded at 04/0001 UTC.


Four-Day Space Weather Forecast Summary

Solar Activity: Low solar activity is expected to continue, with a slight chance of M-class flares.

Solar Wind / Geomagnetic Activity: No CMEs have been assessed as Earth-directed, though there may be one waiting to be analysed. Elevated or Slightly-elevated solar winds are expected to persist until late day 1 into day 2 (4th-5th) before easing as Earth exits into the ambient wind. Geomagnetic activity should remain mainly Quiet with a chance of Unsettled or Active intervals through the period, perhaps increasing to mainly Unsettled with a slight chance of G1/Minor geomagnetic storms on day 3 (6th) with the next Coronal Hole (CH09/+) High Speed Stream arrival, perhaps preceded by a CIR.

Energetic Particles / Solar Radiation: The high energy proton flux (greater than 10 MeV) is expected to remain at background levels. The high energy electron flux (greater than 2 MeV) is expected to remain mostly at background to moderate levels, but with a chance of approaching high levels at the diurnal maxima. The corresponding 24 hour fluence is most expected to remain below the Active threshold (1e8 integrated pfu) but on a steadily increasing trend.

Figure 1. Solar Analysis Valid 03/1900UTC.



Key: Filament ____, Prominence ____, Magnetic Field Line - - -, Polarity +/-, Coronal Holes: Grey shaded area CHxx +/-, Sunspot groups 25xx - Mt Wilson α - β - $\beta\gamma$ - $\beta\gamma\delta$ and Zurich-McIntosh Axx etc.

Geomagnetic Storms:

No Earth directed CMEs are currently expected, though further analysis is required of a possible CME from the earlier C9.2 flare.


Earth is currently embedded within the High Speed Stream of CH07/- giving elevated solar winds. This is a large northward extension of the southern polar crown, spanning nearly 40 degrees of longitude along the southern hemisphere. Solar winds are expected to remain elevated or slightly elevated whilst within this high speed stream (HSS), which is currently expected to persist through to late day 1 into day 2 (4th-5th). The next HSS arrival is expected later on day 3 (6th) from CH09/+.

Geomagnetic activity should remain mainly Quiet with a chance of Unsettled or Active intervals through the period, perhaps increasing to mainly Unsettled with a slight chance of G1 minor geomagnetic storms on day 3 (6th) with the next HSS.

Geo-Magnetic Storm	Level	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC)	Day 2 (00-24 UTC)	Day 3 (00-24 UTC)	Day 4 (00-24 UTC)
Probability (Exceedance)			(%)	(%)	(%)	(%)
Minor or Moderate	G1 to G2	No	5	1	20	10
Strong	G3	No	1	1	1	1
Severe	G4	No	1	1	1	1
Extreme	G5	No	1	1	1	1

Geomagnetic Activity - Earthbound Coronal Mass Ejections- Nil

Figure 2: BGS Estimated planetary Kp




Radio Blackouts - X-Ray Flares:

Solar Activity is currently Low, with only two sunspot regions on the visible disc. The most prominent is in the southeast, AR3068 Esi/Beta-gamma, which has been the source of the two C class flares. AR3071 Hsx/alpha is the other region, situated in the southwest disc. Activity is likely to remain Low, with a slight chance of Moderate activity due to any M-flares that may occur, primarily from AR3068.

X Ray Flares	Level	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC)	Day 2 (00-24 UTC)	Day 3 (00-24 UTC)	Day 4 (00-24 UTC)
Probability			(%)	(%)	(%)	(%)
Active	R1-R2 M Class	No	20	20	20	20
Very Active	R3 to R5 X Class	No	1	1	1	1

Figure 3: GOES16 X-Ray flares



Solar Radiation Storms - (High Energy Protons):

The high energy proton flux (greater than 10 MeV) is currently at Background levels and is expected to remain at this level through the period. AR3068 will need monitoring for any further development, however, with the potential to increase radiation storm risks as it transits across the eastern hemisphere of the solar disc.

Radiation Storms	Level ($\text{cm}^{-2} \text{sr}^{-1} \text{s}^{-1}$)	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC)	Day 2 (00-24 UTC)	Day 3 (00-24 UTC)	Day 4 (00-24 UTC)
Probability (Exceedance)			(%)	(%)	(%)	(%)
Active	$\geq \text{S1}$	No	1	1	1	1
Very Active	$\geq \text{S3}$ *	No	1	1	1	1

* S3 $\geq 10 \text{ MeV} \geq 1000 \text{ pfu}$ and / or $\geq 50 \text{ MeV} \geq 10 \text{ pfu}$. (pfu = $\text{cm}^{-2}\text{sr}^{-1}\text{s}^{-1}$)

High Energy Electrons Event ($\geq 2\text{MeV}$):

The high energy electron flux (greater than 2 MeV) as measured by GOES-16 is initially expected to remain mostly at background levels, but a chance of rising to Moderate and perhaps approaching High (1000 pfu) at diurnal peak. This will most likely occur as solar winds ease later on day 1 into 2 (4th-5th).

The associated 24 hour fluence, however, is expected to remain below the Active threshold (1e8 integrated pfu) throughout. While an increasing trend is likely, electron flux levels are currently not expected to persist at High for a sufficiently long period of time to result in fluence reaching Active levels. This is supported by REFM which is currently giving good guidance.

GEO Electron Environment	Level ($\text{cm}^{-2} \text{sr}^{-1} \text{day}^{-1}$)	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC)	Day 2 (00-24 UTC)	Day 3 (00-24 UTC)	Day 4 (00-24 UTC)
Probability (Exceedance)			(%)	(%)	(%)	(%)
Active	$\geq 2 \text{ MeV}$ $\geq 1 \times 10^8$	No	5	10	10	10
Very Active	$\geq 2 \text{ MeV}$ $\geq 1 \times 10^9$	No	1	1	1	1

Figure 4: MOSWOC REF Model

